



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.


Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

For the given numerical values, $\gamma=30^\circ$, $\beta=60^\circ$, $\lambda-\delta=40^\circ 53' 36''$, we get $\varepsilon=60^\circ$. $\therefore \beta-\varepsilon=60^\circ$. $\therefore \cot x=0$. $\therefore x=90$. Consequently the shadow falls due west.

III. Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Science, Chester High School, Chester, Pa.

Let $OP=100$ feet $=\alpha$ be the tree, AB the meridian, CD' the parallel of latitude, OD the projection of the tree on the plane, OR the shadow, $\angle POD=30^\circ=\gamma$, $\angle DOB=60^\circ=\beta$, $\angle PRD=\text{sun's altitude}=\frac{1}{2}\pi-(\lambda-\delta)$, $\angle RDO=\beta$, $\angle DOE=\frac{1}{2}\pi-\beta$, $\angle PDO=\angle PDR=\angle DEO=\frac{1}{2}\pi$.



$$\therefore PD=\alpha \sin \gamma=50 \text{ feet, } DO=\alpha \cos \gamma=50\sqrt{3}.$$

$$\therefore DO=86.60 \text{ feet, } DE=DO \cos \beta=43.30 \text{ feet, } DR=PD \tan(\lambda-\delta)=\alpha \sin \gamma \tan(\lambda-\delta)=43.30 \text{ feet. } \therefore DE=DR \text{ and } E \text{ and } R \text{ coincide.}$$

\therefore The shadow is due west.

Also solved by EDMUND FISH and A. H. BELL, and H. C. WHITAKER.

80. Proposed by the late SYLVESTER ROBINS, North Branch, N. J.

Exhibit ten initials in that infinite series of integral, rational rhombuses wherein the area of every term is one unit less than the square of its side.

Solution by M. A. GRUBER, A. M., War Department, Washington, D. C.

Let $2a$ and $2b$ be the respective diagonals of a rhombus. As the diagonals of a rhombus bisect each other at right angles, we then have, side of rhombus $=\sqrt{a^2+b^2}$, and area $=2ab$.

$$\therefore \text{From condition of problem, } 2ab=a^2+b^2-1, \text{ or } a^2-2ab+b^2=1.$$

$$\text{Whence } a-b=\pm 1.$$

\therefore The side of rhombus must be the hypotenuse of a right triangle whose legs are consecutive integers.

Several methods for finding successive right triangles of this kind are given in THE AMERICAN MATHEMATICAL MONTHLY, Vol. IV, No. 1, pages 24—27.

Whence we find, for the first ten integral, rational rhombuses, the respective—

Diagonals,		Sides,	and	Areas.
$2a$	$2b$	$\sqrt{a^2+b^2}$		$2ab=a^2+b^2-1$
8.....	6.....	5.....		24
42.....	40.....	29.....		840
240.....	238.....	169.....		28560
1394.....	1392.....	985.....		970224
8120.....	8118.....	5741.....		32959080
47322.....	47320.....	33461.....		1119638520
275808.....	275806.....	195025.....		38034750624
1607522.....	1607520.....	1136689.....		1292061882720
9369320.....	9369318.....	6625109.....		43892069261880
54608394.....	54608392.....	38613965.....		1491038293021224

The general formula for finding sides is $6S_{n-1}-S_{n-2}=S_n$.

Also solved by A. H. BELL, CHAS. C. CROSS, and G. B. M. ZERR.